

Genetic and Biochemical Aspects of Ectoine Biosynthesis in Moderately Halophilic and Halotolerant Methylophilic Bacteria

¹Valentina N. Khmelenina, ¹Ildar I. Mustakhimov, ¹Alexander S. Reshetnikov,

²Marina G. Kalyuzhnaya and ¹Yuri A. Trotsenko

¹Laboratory of Methylophilicity,

G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms,
RAS, Pushchino, Moscow Region, Prospect Nauki 5, Moscow Region, 142290, Russia

²Department of Microbiology, University of Washington, Seattle, WA 98115, USA

Abstract: Problem statement: The cyclic imino acid ectoine is a widely distributed compatible solute synthesized by halophilic and halotolerant bacteria to prevent osmotic stress at high external salinity. This water-keeping compound is used in a variety of commercial cosmetics and therapeutic products. **Approach:** Development of integrated, predictive functional model of the metabolic and regulatory networks of ectoine-producing microbes is an active area of research. In this article we present a brief overview of the current knowledge on genetic and biochemical aspects of ectoine biosynthesis in aerobic halophilic and halotolerant bacteria utilizing C₁ compounds (methylophilic). Although enzymology and genetics of the ectoine biosynthesis in methylophilic bacteria are similar to other halophilic bacteria, the regulatory patterns are different. In all methylophilic bacteria studied, the genes coding for specific enzymes of ectoine biosynthesis: Diaminobutyric Acid (DABA) aminotransferase (EctB), DABA acetyltransferase (EctA) and ectoine synthase (EctC) are organized into ectABC or ectABC-ask, which is linked to gene encoding Aspartokinase isozyme (Ask). **Results:** Remarkably, the methylophilic bacteria possessing a four-gene cluster showed higher halotolerance and accumulated more ectoine than bacteria with a cluster composed of three genes. The DABA acetyltransferases from three methylophilic species have been comparatively characterized. The properties of the enzymes correlate with eco-physiological and metabolic particularities of the host. Some elements of the regulatory system governing the ectoine pathway operation have been revealed in both methane and methanol utilizing bacteria. In *Methylophilium alcaliphilum* transcription of the ectABC-ask operon is initiated from two σ^{70} -like promoters and controlled by the EctR, a MarR-type negative regulator. EctR orthologs were identified in genomes of several heterotrophic halophilic bacteria. Here we present genomic data indicating that similar regulatory system may occur in diverse halophilic and halotolerant bacteria. **Conclusion:** Currently available data suggest that in methylophilic bacteria the ectoine biosynthesis pathways are evolutionary well conserved, particularly with respect to the genes and enzymes involved. However, some differences in the *ect*-gene cluster organization and regulation could be observed.

Key words: Halophilic bacteria, osmoadaptation, ectoine, enzymes, transcriptional regulation

INTRODUCTION

The ability of microbial cells to adapt to fluctuations of external osmolarity (osmoadaptation) via activation of specific mechanisms (osmoregulation) is essential for their survival. Halophilic and halotolerant microorganisms have evolved a number of metabolic, enzymatic and regulatory mechanisms that prevent cell dehydration. Osmoadaptation based on

accumulation of inorganic ions, mostly K⁺ (salt type of osmoadaptation), is employed in extremely halophilic members of *Halobacteriaceae* and *Halanaerobiales*, as well as anaerobic acetogenic species of *Halobacteroides*, *Sporohalobacter*, *Acetohalobium* and sulfate reducers (*Desulfovibrio halophilus*, *Desulfohalobium retbaense*) (Zhilina and Zavarzin, 1990; Galinski and Truper, 1994). Structural and metabolic organization of cells of extremely halophilic

Corresponding Author: Valentina N. Khmelenina, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, RAS, Pushchino, Prospect Nauki 5, Moscow Region, 142290, Russia
Tel: +7 495 925 7448 Fax: +7 495 956 3370

bacteria is well adapted to high intracellular concentrations of ions. Moreover, it often depends on high salinity for optimal operation. For example, malate dehydrogenase of *Halobacterium salinarum* and halophilic archaeon *Haloarcula marismortui* is more acidic and require K^+ for compact folding and enzyme activity (Marhuenda-Egea and Bonete, 2002; Mevarech *et al.*, 2000). Another osmoadaptation strategy is based on synthesis of small organic molecules (osmolytes). It occurs in a majority of bacteria, archaea, fungi, plants and even animals (Galinski and Truper, 1994; da Costa *et al.*, 1998; Ventosa and Nieto, 1995). Osmolytes (also known as compatible solutes) do not interfere with cellular metabolism. They are represented by compounds of different classes: (i) zwitterions (amino acids and their derivatives including ectoines and betaines), (ii) neutral solutes (sugars and polyols) and (iii) anionic solutes where the negative charge is supplied by a carboxylate, phosphate or sulfate (Galinski, 1995; Roberts, 2004; 2005). Many bacteria, including non-halophilic, accumulate compatible solutes at hyperosmotic conditions either through *de novo* synthesis or by uptake from surrounding environment.

Ectoine, a cyclic imino acid (1,4,5,6-tetrahydro-2-methyl-4-pyrimidine carboxylic acid) was originally discovered as a compatible solute in an anoxygenic phototrophs of the *Ectothiorhodospira* group (Galinski *et al.*, 1985). Later, it was found in many aerobic halophilic and halotolerant gram-negative and gram-positive bacteria including *Nocardiopsis*, *Brevibacterium*, *Marinococcus*, *Halomonas*, *Pseudomonas* and *Vibrio* (Ventosa and Nieto, 1995; Galinski, 1995; Kempf and Bremer, 1998; Grant, 2004; Severin *et al.*, 1992). Ectoine and its hydroxylated derivative, hydroxyectoine, are powerful multifunctional bioprotectants that defend against a variety of damaging factors such as heating, freezing, desiccation and UV radiation (Jebbar *et al.*, 1992; Buenger and Driller, 2004; Graf *et al.*, 2008). Both compounds are widely used in medicine and cosmetics as brought-range stabilizers. In spite of a rising commercial interest the genetics and biochemistry of ectoine biosynthesis are intensively studied in various bacteria. We refer here to the genetic and enzymatic aspects of the ectoine biosynthesis pathway in aerobic methylotrophic bacteria which utilize single carbon compounds (methane, methanol or methylamines) as the carbon and energy sources. These moderately halophilic and halotolerant methylotrophs have been isolated from soda lakes and (hyper) saline environments. The members of the genera *Methylomicrobium*, *Methylobacter*, *Methylophaga*

(*Gammaproteobacteria*) were shown to synthesize ectoine as an osmoprotectant along with sucrose and glutamate (Khmelenina *et al.*, 1999; Doronina *et al.*, 2003a; 2003b; Kalyuzhnaya *et al.*, 2001; 2008). Ectoine was also identified in cells of *Methyloarcula* species (*Alphaproteobacteria*) grown at high salinity (Doronina *et al.*, 2000).

MATERIALS AND METHODS

Genetic aspects of ectoine biosynthesis: The ectoine biosynthesis represents a branch of the metabolic road for aspartate family amino acids synthesis (Fig. 1) and involves three special enzymes: Diaminobutyric Acid (DABA) aminotransferase (EctB) that catalyses amination of aspartate semialdehyde into diaminobutyric acid, DABA acetyltransferase (EctA) acetylating DABA into $N\gamma$ -acetyl diaminobutyric acid and ectoine synthase (EctC) that forms ectoine by cycling of $N\gamma$ -acetyl diaminobutyric acid (Peters *et al.*, 1990). In various halophilic bacteria studied, the genes coding for the enzymes form a cluster ectABC (Canovas *et al.*, 1998; Louis and Galinski, 1997; Goller *et al.*, 1998; Kuhlmann and Bremer, 2002). Some bacteria possess an additional enzyme that converts ectoine to hydroxyectoine (ectoine hydroxylase, encoded by ectD) (Canovas *et al.*, 1999; Prabhu *et al.*, 2004; Bursy *et al.*, 2007). The ect-genes are usually, but not always, organized into more or less compact operon.

Analysis of complete microbial genomes showed that the ect-genes are widespread among *Proteobacteria* and *Actinobacteria* (Fig. 2). The genetic signature of the ectoine pathway, an *ectABC* operon, was also identified in genome of *Nitrosopumilus maritimus*, a non-thermophilic archaeon isolated from a salt-water aquarium. Rather high homology of the ectoine biosynthetic genes in microorganisms of different taxonomic position and physiology is a reliable indication of the evolutionary conservation of the biochemical pathway (Kuhlmann and Bremer, 2002).

High homology of the *ect*-genes in microorganisms allowed identification of the genes in some aerobic methylotrophs by using PCR-surveys (Reshetnikov *et al.*, 2006). *ectABC* or *ectABC*-ask clusters were identified in six species of genera *Methylomicrobium*, *Methylobacter*, *Methylophaga* and *Methyloarcula*. The organization of the *ect*-genes (three or four gene cluster) correlates with halotolerance of the host strain. Two halotolerant methanotrophs, *Methylomicrobium kenyense* AMO1 and *Methylobacter marinus* 7°C possess three gene operon (*ectABC*), accumulate ectoine up to 70 mg g⁻¹ of Dry Cell Weight (DCW) and are capable of growth at salinity 4-5% NaCl.




Fig. 1: Pathway of ectoine biosynthesis from aspartic acid

In contrast, methanotrophic species *Methylomicrobium alcaliphilum* and methanol- and methylamine-utilizers *Methylophaga alcalica*, *Methylophaga thalassica* and *Methylarcularia marina* possess ectABC-ask operon, can grow at two-fold higher salinity (10-12% NaCl) and accumulate >20 mg of ectoine g⁻¹ of DCW. This implies an important role of the ectoine pathway specific aspartokinase in some methylotrophic species. It was demonstrated that in *Mm. alcaliphilum*, genes ectABC and ask are co-transcribed and thus the aspartokinase must be osmotically controlled (Reshetnikov *et al.*, 2006). One could speculate that the occurrence of the aspartokinase isozyme may provide an advantage at high osmolarity as it makes ectoine biosynthesis independent from complex machinery regulating the amino acid biosynthesis. However, in this case a culture must possess an additional enzyme that could fulfill the pool of amino acid for protein synthesis at low salinity. Indeed, a copy of gene coding for aspartokinase with 77% identity of the translated amino acids is present in *Mm. alcaliphilum* as followed from a draft genome sequence analyses (<https://www.genoscope.cns.fr/agc/mage>).

Analyses of the publicly available complete microbial genomes showed that at least 30 bacterial

species harbor homologs of the ectABC-ask genes. Bacteria possessing an ask gene immediately downstream of ectABC are exclusively gram-negative. In the majority of the proteobacterial halophiles, a second, additional copy of the aspartokinase gene was identified. Two and three additional ask genes not linked to the ect-gene cluster occur in *Photobacterium profundum* and *Vibrio cholerae*, respectively. Interestingly, no aspartokinase gene outside of the ect-cluster was found in three species of marine bacteria, *Oceanobacter* sp. RED65, *Sphingomonas* sp. SKA58 and *Lentisphaera araneosa*. In these bacteria the sole ectoine-associated Ask should support both cellular processes-osmoresistance and protein biosynthetic needs. Most likely in these marine bacteria the ect-operon and linked ask gene are constitutively expressed. However, this hypothesis remains to be tested by the comprehensive description of the gene expression and the enzyme properties.

It should be mentioned that in gram-positive bacteria ectoine is not the sole osmoprotectant and other organic compatible solutes belonging to the glutamate family aminoacids, glutamate, glutamine and/or proline contribute to osmotic balance (Kuhlmann and Bremer, 2002; Lo *et al.*, 2009; Saum and Muller, 2007; 2008a; 2008b).




Fig. 2: Phylogenetic tree based on deduced amino acid sequences from *ectB* genes of methylotrophs (underlined) and those from other bacteria possessing *ect* genes. The tree was constructed by minimal evolution method. In bracket the accession numbers of the respective genes and whole genomes presented in Database

This may require a complex regulation since the carbon flow must be also directed to biosynthesis of the osmoprotectants not derived from aspartic acid. It may explain the absence of the *ask* homologues in the *ect*-gene cluster of gram-positive bacteria. Remarkably, the halophilic *Firmicutes* typically possess multiple *ask* homologues not linked to the *ect*-genes (Lo *et al.*, 2009).

Phylogenetic relationships of the EctB from bacteria of different physiological groups are shown in Fig. 2. The topology of the EctB tree was similar to those of EctA- and EctC-based phylogenetic trees (data not shown). In general, the Ect proteins from bacteria belonging to the Alpha- and *Gammaproteobacteria*, *Firmicutes* and *Actinobacteria* comprise separate

branches on the trees. This may imply an ancient acquisition of the ect-genes from common ancestor and prolonged evolution inside the respective microbial phylum. Interestingly, the Ect proteins of methane- and methanol-utilizing bacteria of the *Methylobacterium* and *Methylophaga* genera are combined in the coherent group on the tree being most closely related to other representatives of the *Gammaproteobacteria*. On the other hand, the Ect proteins from *Mm. kenyaense* AMO1 are more identical to those of the methanol-utilizers than other methanotrophs. Since the bacterium carries the ectABC gene cluster, a recent loss of the ask gene may be proposed. Almost identical Ect proteins (94.8, 98.2 and 98.5% for EctA, EctB and EctC) are present in two strains of the *Mm. alcaliphilum*, 20Z and ML1, isolated from the Tuva soda lake (Russia) and Mono Lake (USA), respectively. Hence, geographical distance of the bacterial habitats has not resulted in significant divergence of the ect genes.

Notably, Ect proteins from *Mb. marinus* 7°C, a methanotroph belonging to the *Gammaproteobacteria*, are only distantly related to ectoine biosynthetic enzymes of other methylotrophs (only 37-51% identities to EctABC sequences of *Mm. alcaliphilum* 20Z) and fall with proteins of the marine representatives of *Proteobacteria*. Ectoine biosynthesis proteins from *Mb. marinus* 7°C are closely related to those from the marine *Mariprofundus ferrooxydans* (*Zetaproteobacteria*) (55-80% identities). However, in *M. ferrooxydans* the ectoine biosynthetic pathway is encoded by an ectABC-ask operon, while the ect-cluster from strain 7C lacks ask gene. So far, *Mb. marinus* is the only representative of the *Methylobacter* genus that is salt-resistant and synthesizes ectoine. Other described species of the genus are non-halophilic. It could be proposed that the ect-operon ubiquity in ancient prokaryotic world that was largely marine, followed by loss in lineages that became adapted to a terrestrial environment (Lo *et al.*, 2009). It is also possible that an ancestor, terrestrial species of *Methylobacter* adapted to marine ecosystem by acquiring ectoine biosynthesis genes from phylogenetically distant species.

The biosynthesis of hydroxyectoine proceeds via direct hydroxylation of ectoine catalyzed by ectoine hydroxylase (EctD) (Bursy *et al.*, 2007; Reuter *et al.*, 2010). In *Mm. alcaliphilum*, an orf with a high homology to ectD (40-42% identity to ectoine hydroxylases from *Streptomyces avermitilis* MA-4680 and *S. chrysomallus*) was identified 360 bp downstream of the ectABC-ask operon (Reshetnikov *et al.*, 2006). No hydroxyectoine accumulation was detected in *Mm. alcaliphilum* cells grown in the presence of 3 or 9%

NaCl. However, it does not exclude that the methanotroph can produce the bioprotectant at some environmental perturbations. NR-Database searches revealed that genomes of 67 proteobacteria and one archaeon, *Nitrosopumilus maritimus*, contain homologues of the *ectD* gene (with identities ranging from 41-55% to that of *V. salexigens*) (Reuter *et al.*, 2010). *ectD*-like gene is either a part of the ect-cluster or it locates separately. Majority of the microbial genomes contain only one copy of the *ectD* gene, with exception to *Rhodococcus opacus* B4, *Marinobacter aquaeolei* VT8 and *C. salexigens*, that possess two types of *EctD*-proteins (Reuter *et al.*, 2010). In the case of *C. salexigens*, only one of the *EctD*-like enzymes contributes to the production of 5-hydroxyectoine (Garcia-Esteva *et al.*, 2006) and the *ectD* gene is essential for thermoprotection of the bacterium.

It should be mentioned that an alternative pathway for transformation of N γ -acetyl DABA into hydroxyectoine via 3-hydroxy-N γ -acetyl DABA without ectoine formation step was proposed but this was not enzymatically proved (Canovas *et al.*, 1999).

RESULTS AND DISCUSSION

Some properties of the ectoine biosynthetic enzymes:

In the conditions of hyperosmotic stress (from 0.5-1 M NaCl), *H. elongata* cells with blocked by chloramphenicol protein biosynthesis, accumulated ectoine at the same level as the cells with active protein synthesis. Thus, the regulation of ectoine synthesis can proceed at the enzyme level (Kraegeloh and Kunte, 2002). Three ectoine biosynthesis specific enzymes have been purified from *H. elongata* DSM2581 (Ono *et al.*, 1999). The enzymes are characterized by similar parameters for the maximal activity (pH 8.2-9.0, t = 15-20°C and 0.4-0.5 M NaCl). Since the optimal salt concentrations for the enzymes were lower than those in the medium supporting maximal growth rate of the bacteria, relatively low intracellular concentrations of ions should be maintained (Kraegeloh and Kunte, 2002). Indeed, Na⁺ content of 0.04-0.2 M was found in cells of *Vibrio costicola* и *Brevibacterium* sp. growing at high NaCl concentrations (Gilboa *et al.*, 1991; Nagata *et al.*, 1995).

DABA aminotransferase (EctB, EC 2.6.1.76): DABA aminotransferase from *H. elongata* is a homo-hexameric (~250 kDa) pyridoxal phosphate-dependent enzyme that requires K⁺ for activity and stability. The enzyme is more active in the presence of 0.01-0.5 M KCl, than in the presence of NaCl (Ono *et al.*, 1999). K⁺ requirement is inherent to many enzymes from extremely halophilic

eubacteria and archae with the salt-type osmoadaptation (Marhuenda-Egea and Bonete, 2002; Mevarech *et al.*, 2000; Toney *et al.*, 1995). DABA aminotransferase is specific to L-glutamate, as an amino donor (K_m 9.1 mM) and to D, L-aspartyl semialdehyde (K_m 4.5 mM). The reaction catalyzed by DABA aminotransferase was predicted to be a limiting step of the ectoine biosynthesis pathway thus explaining the absence of DABA in cells of *H. elongata* KS3 (Ono *et al.*, 1999). Two putative DABA aminotransferase genes, one in ectABC-ask operon and another in a cluster of genes presumably involved in ectoine degradation, were found in the genome of *M. alcaliphilum*.

Interestingly, DABA aminotransferase was detected in several bacteria that do not synthesize ectoine. In *Acinetobacter baumannii*, DABA aminotransferase is involved in biosynthesis of diamonopropane, a component of cell wall peptidoglycan (Ikai and Yamamoto, 1997). In *Paenibacillus polymyxa* the enzyme is a part of the polymyxin, a peptide antibiotic, formation (Ono *et al.*, 1999). In *A. baumannii*, it is specific to L-glutamate, whereas in *Xanthomonas* species to L-alanine as the amino donor (Rao *et al.*, 1969).

DABA acetyltransferase (EctA, EC 2.3.1.178): DABA acetyltransferase from *H. elongata* was purified (≈ 400 fold) and was only partially characterized due to low stability of the enzyme (Ono *et al.*, 1999). His-tagged DABA acetyltransferases were purified from *Mm. alcaliphilum* 20Z, *M. alcalica* and *M. thalassica* (Reshetnikov *et al.*, 2005; Mustakhimov *et al.*, 2008). The enzyme from these methylotrophs existed as a homodimer with the subunit molecular mass of ~ 20 kDa and had no requirement for divalent ions.

Some differences in the enzyme properties that correlated with eco-physiologies of these bacteria were described. Thus, the DABA acetyltransferase from the neutrophilic *H. elongata* and *M. thalassica* was more active at lower pH (pH 8.2 or pH 9.0), than that from alkaliphilic species *Mm. alcaliphilum* and *M. alcalica* (pH optima ≥ 9.5). Unlike enzymes from *M. alcalica* and *Mm. alcaliphilum*, DABA acetyltransferase from the marine bacterium *M. thalassica* was considerably inhibited by carbonates (Mustakhimov *et al.*, 2008). Bearing in mind that growth of the former alkaliphilic methylotrophs is stimulated by carbonates, this feature of the enzyme corresponds to *in situ* surroundings of the strains. Interestingly, 1 mM Cu^{2+} completely inhibited activity of the DABA acetyltransferase from *M. alcalica* and 47% that of *M. thalassica*. In contrast, no inhibitory effect of Cu^{2+} was found for the enzyme from methanotroph *Mm. alcaliphilum* and this also

correlates with an important role of copper in methane oxidation in the culture, since Cu is an essential component of the particulate methane monooxygenase (Murrell *et al.*, 2000).

Another intriguing feature of DABA acetyltransferases from methylotrophic bacteria was the effect of ionic strength on enzyme activity. Like the enzyme from *H. elongata*, DABA acetyltransferase from methanotroph *Mm. alcaliphilum* was activated by 0.2 M NaCl implying the "halophilic nature" of the protein. Contrary, the enzymes from methylotrophic cultures, *M. thalassica* and *M. alcalica* were inhibited by salts. We may speculate that methanol is more effective substrate for supporting ion extrusion mechanisms in comparison to methane or glucose. As a result, low concentrations of monovalent inorganic ions could be maintained in cytoplasm of the methanol-utilizing bacteria eliminating need for ectoine biosynthetic enzymes that are adapted to high ionic strength.

Ectoine synthase (EctC, EC 4.2.1.108): The homogenous ectoine synthase from *H. elongata* was purified in the presence of 1 mM N γ -acetyl-DABA and 2 M NaCl as stabilizing compounds. The molecular mass of the native enzyme remains unclear due to loss of the enzyme activity after gel-filtration at 0.5 M NaCl. The EctC-protein contains enhanced levels of aspartate and glutamate. The enzyme is specific to N γ -acetyl-DABA, however the N-acetyl group in α -position could not be involved in the cycling process (Ono *et al.*, 1999). The recombinant ectoine synthase was purified from *Mm. alcaliphilum* 20Z with activity ~ 64 U mg^{-1} . The protein is a 35 kDa homodimer.

Ectoine hydroxylase (EctD, EC 1.14.11): Was purified from the moderate halophile *Virgibacillus (Salibacillus) salexigens* (Bursy *et al.*, 2007) and from the *Streptomyces coelicolor* (Bursy *et al.*, 2008). EctD from *V. salexigens* is a monomeric protein of molecular mass 34 kDa being a member of the non-heme iron (II)- and 2-oxoglutarate-dependent dioxygenases. The reaction depends on iron (II), molecular oxygen and 2-oxoglutarate. The similar properties were observed for enzyme from *Streptomyces coelicolor* with respect to optimal pH (pH 7.5), temperature (32°C), K_m values for ectoine (3.5 and 2.6 mM) and co-substrate 2-oxoglutarate (5.2 and 6.2 mM). Contrary to preferential production of 5-hydroxyectoine by *S. coelicolor* grown at 39°C (Bursy *et al.*, 2007), the high temperature optimum of the EctD enzyme implies that there is no specific thermoactivated regulation of the hydroxylase.

The crystal structure analysis of the *V. salexigens* EctD (Reuter *et al.*, 2010) showed that the folding of the protein is similar to the human phytanoyl-CoA 2-hydroxylase (McDonough *et al.*, 2005). The core of the EctD structure consists of a double-stranded β -helix forming the main portion of the active site of the enzyme. The positioning of the iron ligand in the active site of EctD is mediated by an evolutionarily conserved 2-His-1-carboxylate iron-binding motif. The side chains of the three residues forming this iron-binding site protrude into a deep cavity in the EctD structure that also harbors the 2-oxoglutarate binding site. Despite high homology and similar reactions catalyzed by the ectoine hydroxylase and L-proline hydroxylase (Hausinger, 2004), the EctD from *S. coelicolor* had no additional L-proline hydroxylase activity (Bursy *et al.*, 2007).

Transcriptional regulation of the ectoine biosynthesis genes:

Regulation of the ectoine biosynthesis genes in heterotrophic bacteria: To date, there are only fragmentary data on how an external salinity regulates expression of ectoine biosynthetic genes. In *Bacillus pasteurii* (Kuhlmann and Bremer, 2002), *Marinococcus halophilus* (Bestvater and Galinski, 2002), *Halobacillus halophilus* (Saum and Muller, 2008b) and *S. salexigens* (Bursy *et al.*, 2007), transcription of the *ectABC-ask* or *ectABC* genes proceeded as polycistronic mRNAs and subjected to osmotic conditions. The osmoregulated expression of *ectABC* was also revealed in gram-negative *Brevibacterium epidermis* (Onraedt *et al.*, 2004), *Mm alcaliphilum* (Reshetnikov *et al.*, 2006; Mustakhimov *et al.*, 2010) and *Chromohalobacter salexigens* (Calderon *et al.*, 2004).

In *C. salexigens*, the *ectABC* operon is transcribed from several promoter regions. Four putative promoters (PectA1-4) are located upstream of the *ectA* gene and one internal promoter PectB is upstream of the *ectB* gene (Calderon *et al.*, 2004). The consensus sequences of the *ect* promoters closely resembled those of σ^{70} - (PectA1 and PectA2), σ^S - (PectA3) and σ^{32} - (PectB) dependent promoters of *E. coli*. Both PectA and PectB promoter regions were shown to be osmoregulated. Transcription from the PectB was also enhanced upon increasing growth temperature. Expression of the reporter gene *lacZ* under control of the PectA in recombinant *C. salexigens* and *E. coli* cells was considerably increased when the cultures approached stationary growth phase. Moreover, in the mutant *E. coli* lacking the *rpoS* gene, the expression of the PectA-*lacZ* fusion was much lower than in the wild type strain. This was consistent with maximal ectoine

accumulation in the stationary growth phase (Canovas *et al.*, 1999). It could be reasonable to propose an involvement of the general stress factor σ^S , however, no *rpoS*-like gene has been identified in *C. salexigens* (Calderon *et al.*, 2004). Participation of different transcriptional sigma factors in expression of the *ect*-genes was not confirmed by an appropriate genetic study. Supplementation of the growth media with ectoine or glycine betaine decreased the transcription of PectA-*lacZ* and PectB-*lacZ* (Calderon *et al.*, 2004). Altogether with constitutive *ect*-genes transcription from PectA and PectB at low osmolarity, this implies an involvement of other transcriptional factors.

The recent studies have shown a connection between the iron homeostasis and the osmoreponse in *C. salexigens* (Argandona *et al.*, 2010). An iron homeostasis regulator, Fur, has been described as a potential regulatory link between salinity and iron metabolism. Corresponding six-gene operon *cfuABC-fur-hisI-orf6* was identified downstream of the *ectABC* genes. Fur boxes were found in promoters of the *cfuABC-fur-hisI-orf6* and *ectABC* operons. It was shown that Fur mediates the osmoregulated inhibition by iron of *cfuABC-fur-hisI-orf6* expression and functioned as a positive regulator of the *ectABC* genes under high-salinity conditions.

In gram-positive bacteria, ectoine biosynthesis also seems to be triggered by an osmotic stress. The vegetative σ^A -dependent promoter (equivalent to σ^{70} -like promoter of gram-negative bacteria) was found upstream of the *ectABC* in *Bacillus subtilis*, *B. pasteurii* and *S. salexigens* (Kuhlmann and Bremer, 2002; Bursy *et al.*, 2007). In *Marinococcus halophilus*, the transcription of *ectABC* genes was initiated from three individual σ^{70}/σ^A -promoters located upstream of each gene (Bestvater and Galinski, 2002). The σ^{70}/σ^A -promoter sequences were amplified and cloned upstream of *gfp* in vector pBR322. A linear increase in fluorescence of *E. coli* cells upon increase of NaCl concentration in the medium was detected, however, the addition of ectoine and betaine decreased the fluorescence signal (Bestvater and Galinski, 2002).

Moderately halophilic *Halobacillus halophilus* served as a prominent model bacterium to decipher regulatory mechanisms of the osmolytes biosynthesis. When grown at moderate salinity (1 M NaCl), *H. halophilus* accumulated glutamate and glutamine as the major compatible solutes whereas ectoine and proline were predominantly produced at very high salinities (Saum and Muller, 2008a; 2008b). Ectoine/proline ratio that was low in cells growing exponentially at different salinities (2 and 3 M NaCl) enhanced more than 1000-fold in the stationary growth phase. The proline

exhausting in the stationary phase cells (the amino acid proline can be also used as the carbon or energy source) may be the major reason for the observed growth phase-dependent switches (Saum and Muller, 2008a; 2008b).

Osmotic up-shock of *H. halophilus* cells led to the enhanced levels of mRNAs of ectABC and gdh1, glnA2 and proH genes encoding respectively glutamate dehydrogenase, glutamine synthetase and pyrroline-5-carboxylate reductase.. The expression of ect-genes reached the maximal level when concentration of transcripts for genes for glutamate, glutamine and proline biosynthesis returned almost to the initial levels. The ectABC transcript concentration was highest in the presence of NaNO₃. Na-gluconate was as efficient as NaCl while Na-glutamate led only to a minor increase of the transcripts. Hence, transcription of the ect-genes did not necessarily depend on the presence of chloride anions (Saum and Muller, 2008b).

It was demonstrated that in *C. salexigens* and *Streptomyces griseus*, intracellular levels of ectoine increase in response to high growth temperature (Calderon *et al.*, 2004; Malin and Lapidot, 1996). These findings suggest a role of ectoine as a protectant against the detrimental effects of high temperatures. In contrast, the ectoine biosynthesis in *V. pantothenicus* was triggered by low growth temperatures (Kuhlmann *et al.*, 2008). The highest level of ectoine and the ectABC transcripts were detected *V. pantothenicus* cells subjected simultaneously to chilling (15°C) and salt stress (0.7 M NaCl).




Fig. 3: Involvement of the EctR in transcriptional regulation of ectoine biosynthesis. At the low osmolarity, ectABC-ask genes of *Mm. alcaliphilum* are constitutively transcribed from weak promoter ectA2p. EctR1 sterically inhibits binding of the RNA polymerase with -10 sequence of promoter ectA1p repressing transcription from osmoregulating promoter ectA1p. At the increasing of medium salinity, EctR-DNA complex is dissociated making promoter ectA1p to be accessible for RNA polymerase

Several additional regulatory elements of hydroxyectoine biosynthesis can be recognized in *C. salexigens* and *S. salexigens*. In the latter, mRNA transcribed from the ectD promoter was 900 b.p. being larger than ectD sequence. An orf similar to MarR-family of transcriptional regulators was revealed downstream of the ectD (Bursy *et al.*, 2007). However, the putative regulatory protein is disrupted by two stop codons and no start codon could be found. In *C. salexigens*, the orf (gene ectR) was found upstream of ectD (Vargas *et al.*, 2008; Garcia-Esteva *et al.*, 2006). Deletion of the ectR gene resulted in decreasing of hydroxyectoine level at high salinity and high temperature, in comparison to the wild type cells. Thus, EctR (presumably belonging to the AraC family of transcriptional regulators) could be an activator of the ectD transcription.

Recent findings in transcriptional regulation of ectoine biosynthesis genes in methylotrophs: In *M. alcaliphilum*, transcription of the ectABC-ask operon is initiated from two σ^{70} -dependent promoters ectAp1 and ectAp2 (Fig. 3). The predicted -10 (TACTAT) and -35 (TGGACA) regions of ectAp1 showed rather high level of identity with the consensus sequence of the *E. coli* σ^{70} -recognized promoter (Reshetnikov *et al.*, 2006; Mustakhimov *et al.*, 2010). The putative -10 and -35 sequences of the ectAp2 promoter differ from the respective regions of the *E. coli* σ^{70} -promoter thus suggesting that expression from ectAp2 may be less effective, than from ectAp1.

Upstream of the ectA gene an orf (gene ectR1) encoding the MarR-like transcriptional regulator with 12-20% identity of translated amino acid sequences was found. Despite low identity, the protein has a structure analogous to the MarR-family regulators: it has the Helix-Turn-Helix (HTH) DNA-binding motif flanking the "wing 1" region (Mustakhimov *et al.*, 2010; Hong *et al.*, 2005; Wilkinson and Grove, 2006). Regulatory function of EctR1 in ectoine biosynthesis was elucidated by characterization of the ectR1 knockout mutant. A promoter-reported system (ectP-gfp fusion) was constructed and introduced into *Mm. alcaliphilum* wild type and mutant strains. The fluorescence signal in the strain lacking ectR1 was 2-3 folds higher compared to wild type cells. The DABA acetyltransferase activity was also 2-6 times higher in the mutant cells. Thus, EctR1 negatively controlled transcription of the ect-genes in *Mm. alcaliphilum*. Expression of the ectABC-ask operon in the ectR1 mutant was still activated by increasing of the medium salinity (from 1-6% NaCl) thus indicating that *Mm. alcaliphilum* may possess several regulatory systems.




Fig. 4: Phylogenetic tree of putative transcriptional regulators EctR of halophilic bacteria and other regulators of the MarR-family

Transcription of the gene *ectR1*, in turn, was carried out from a single σ^{70} -like promoter (Fig. 3). The promoter region of *ectR1*, *ectR1p*, is located between *ectAp1* and *ectAp2*, suggesting, that its transcription may be controlled by the EctR. The autoregulation may be described for some other MarR proteins. The purified EctR1 specifically binds to the promoter region of *ectABC*-ask operon (Mustakhimov *et al.*, 2010). The EctR1 binding site contains a pseudopalindromic sequence (TATTTAGT-GT-ACTATATA) composed of 8-bp half-sites separated by 2 bp suggesting dimeric association of the EctR1 with the DNA where each protein subunit binds with an inverted repeat (Mustakhimov *et al.*, 2010). Indeed, gel-filtration studies showed that EctR is a dimer both in free solution (m.m. 44-45 kDa) and in DNA binding state (m.m. 50-55 kDa). Hence, it may be proposed that at

the low osmolarity, the *ectABC*-ask genes of *Mm. alcaliphilum* are constitutively transcribed from the weak promoter *ectA2p* (Fig. 3). EctR1 sterically inhibits the RNA polymerase binding with -10 sequence of the promoter *ectA1p* and thus represses transcription from the osmoregulating promoter *ectA1p*. At the elevated external salinity, EctR-DNA complex dissociates making the promoter *ectA1p* to be accessible for RNA polymerase (Fig. 3).

The EctR1 orthologs were identified in other halophilic bacteria. Our analysis of the DNA fragment containing the ectoine biosynthetic genes in the methanol-utilizing bacterium *M. alcalica* (Mustakhimov *et al.*, 2009) showed the presence of an orf with high homology to the *ectR1* gene from *Mm. alcaliphilum* (73% identity of translated amino acids). Moreover, a simple NCBI Blast search revealed several

ectR1-like genes located immediately upstream of the ectoine gene cluster in 17 halophilic bacterial species (Fig. 4). Between them, the orfs of *Oceanospirillum* sp. (EAR60187), *Nitrosococcus oceani* (ABA57535), *Saccharophagus degradans* (ABD80450), *Reinekea* sp. (ZP_01114878) and *Oceanobacter* sp. (EAT11341) showed the highest identities of translated amino acid sequences with the EctR1 from *Mm. alcaliphilum* (35.5, 42.2, 45.6, 51.7 and 55.1%, respectively) which are higher than with other MarR-family transcriptional proteins (<20%). We propose that putative EctR1 of these halophiles represents a separate subfamily in the MarR-family transcriptional proteins (Fig. 4). These results suggest that EctR-mediated regulatory system controlling ectoine biosynthesis at the transcriptional level may operate in diverse halophilic and halotolerant bacteria.

So far, it is not clear how changes in external osmolarity may regulate expression of the ectoine biosynthetic genes via the EctR1. It seems logical to propose that DNA binding ability depends on the external salinity. However, the activity of EctR1 could not be directly regulated by salt, since the expression of the ectABC-ask in *Mm. alcaliphilum* 20Z proceeds at low osmolarity. In any event, the tight relationship between salinity level and ectoine biosynthesis implies an involvement of complex multi tiered regulatory network, including inducers, sigma factors, transcriptional repressors and activators.

CONCLUSION

The necessity of unraveling the principles of organization and regulation of the genes and enzymes involved in ectoine biosynthesis by aerobic methylotrophs is conditioned by practical demands for a technology for production of the multifunctional valuable bioprotectants from cheap sources of carbon, such as methane and methanol. Large-scale production of ectoine has been achieved by the "Bitop" and "Merck" (Germany) using heterotrophic bacterium *Halomonas elongata*. This milking bioprocess is based on the use of the glucose, L-amino acids (glutamate) and highly saline medium (12% NaCl) (Sauer and Galinski, 1998). Aerobic moderately halophilic and halotolerant methylotrophic bacteria when grown at 9% NaCl are capable of accumulation ectoine up to 20% of DCW and therefore, are promising ectoine producers from renewable methane and/or methanol. The differences in ectoine accumulation may be caused by genetically defined regulatory mechanisms in the bacterial producers. Although the enzymology and genetics of the ectoine biosynthesis pathway in

methylotrophs are similar to halophilic bacteria, they may have quite different regulatory patterns. For instance, the presence of the osmotically controlled aspartokinase in methylotrophs could make ectoine synthesis rather independent on other amino acids biosynthesis. Such co-ordination of aspartylphosphate synthesis, a common precursor of both ectoine and the amino acids, may provide an advantage during methylotrophic growth at high external osmolarity.

Osmoadaptation of aerobic methylotrophs, besides the osmoprotective compatible solutes biosynthesis (ectoine, glutamate and in some case sucrose), includes other structure-functional mechanisms such as changes in phospholipids fatty acids composition and in bioenergetic machinery. To date, it is not possible to describe whole regulatory cascade from sensing signals and receivers on a cell membrane to the real metabolic and structural rearrangements. Clarifying the nature of initial signals and sensors on a cell membrane and signal transduction to potential transcriptional regulators is a challenging task for future genomic transcriptomic and proteomic studies. The elucidation of cell responses to external perturbations, such as temperature, pH and others will shed more light on still enigmatic mechanisms of cross-adaptation of bacteria to fluctuating environmental conditions.

ACKNOWLEDGMENT

This study was supported by grants from the Russian Foundation for Basic Research (10-04-01224-a; 09-04-92520-ИК_a), the Ministry of Education and Science RF (RNP2.1.1/605), Russian State Contract 02.740.11.0296, CRDF Rub1-2946-PU-09 and the U.S. National Science Foundation (MCB-0842686). The authors are gratefully acknowledged to the anonymous referees for useful comments and advices.

REFERENCES

- Argandona, M, J.J. Nieto, F. Iglesias-Guerra, M.I. Calderon and R. Garcia-Esteba *et al.*, 2010. Interplay between iron homeostasis and the osmotic stress response in the halophilic bacterium *Chromohalobacter salexigens*. *Applied Environ. Microbiol.*, 76: 3575-3589. DOI: 10.1128/AEM.03136-09
- Bestvater, T. and E.A. Galinski, 2002. Investigation into a stress-inducible promoter region from *Marinococcus halophilus* using green fluorescent protein. *Extremophiles*, 6: 15-20. PMID: 11878557
- Buenger, J. and H. Driller, 2004. Ectoin: An effective natural substance to prevent UVA-induced premature photoaging. *Skin Pharmacol. Physiol.*, 17: 232-237. DOI: 10.1159/000080216

- Bursy, J., A.J. Pierik, N. Pica and E. Bremer, 2007. Osmotically induced synthesis of the compatible solute hydroxyectoine is mediated by an evolutionarily conserved ectoine hydroxylase. *J. Biol. Chem.*, 282: 31147-31155. DOI: 10.1074/jbc.M704023200
- Bursy, J., A.U. Kuhlmann, M. Pittelkow, H. Hartmann and M. Jebbar *et al.*, 2008. Synthesis and uptake of the compatible solutes ectoine and 5-hydroxyectoine by *Streptomyces coelicolor* A3(2) in response to salt and heat stresses. *Applied Environ. Microbiol.*, 74: 7286-7296. DOI: 10.1128/AEM.00768-08
- Calderon, M.I., C. Vargas, F. Rojo, F. Iglesias-Guerra and L.N. Csonka *et al.*, 2004. Complex regulation of the synthesis of the compatible solute ectoine in the halophilic bacterium *Chromohalobacter salexigens* DSM 3043^T. *Microbiology*, 150: 3051-3063. DOI: 10.1099/mic.0.27122-0
- Canovas, D., C. Vargas, M.I. Calderon, A. Ventosa and J.J. Nieto, 1998. Characterization of the genes for the biosynthesis of the compatible solute ectoine in the moderately halophilic bacterium *Halomonas elongata* DSM 3043. *Syst. Applied Microbiol.*, 21: 487-497. PMID: 9924816
- Canovas, D., N. Borges, C. Vargas, A. Ventosa and J.J. Nieto *et al.*, 1999. Role of N γ -acetyldiaminobutyrate as an enzyme stabilizer and an intermediate in the biosynthesis of hydroxyectoine. *Applied Environ. Microbiol.*, 65: 3774-3779. PMID: 10473374
- Da Costa, M.S., H. Santos and E.A. Galinski, 1998. An overview of the role and diversity of compatible solutes in bacteria and archaea. *Adv. Biochem. Eng. Biotechnol.*, 61: 117-153. PMID: 9670799
- Doronina, N.V., Y.A. Trotsenko and T.P. Tourova, 2000. *Methylophaga marina* gen. nov., sp. nov. and *Methylophaga terricola* sp. nov.: Novel aerobic, moderately halophilic, facultatively methylotrophic bacteria from coastal saline environments. *Int. J. Syst. Evol. Microbiol.*, 50: 1849-1859. PMID: 11034496
- Doronina, N.V., T.D. Darmaeva and Y.A. Trotsenko, 2003a. *Methylophaga alcalica* sp. nov., a novel alkaliphilic and moderately halophilic, obligately methylotrophic bacterium from an East Mongolian saline soda lake. *Int. J. Syst. Evol. Microbiol.*, 53: 223-229. DOI: 10.1099/ijs.0.02267-0
- Doronina, N.V., T.D. Darmaeva and Y.A. Trotsenko, 2003b. *Methylophaga natronica* sp. nov., a new alkaliphilic and moderately halophilic, restricted-facultatively methylotrophic bacterium from soda lake of the Southern Transbaikalian region. *Syst. Applied Microbiol.*, 26: 382-389. PMID: 14529181
- Galinski, E.A., 1995. Osmoadaptation in bacteria. *Adv. Microbiol. Physiol.*, 37: 273-328. PMID: 8540423
- Galinski, E.A. and H.G. Truper, 1994. Microbial behavior in salt-stressed ecosystems. *FEMS Microbiol. Rev.*, 15: 95-108. DOI: 10.1111/j.1574-6976.1994.tb00128.x
- Galinski, E.A., H.P. Pfeiffer and H.G. Truper, 1985. 1, 4,5,6, -Tetrahydro-2-methyl-4-pyrimidinecarboxylic acid, a novel cyclic acid from halophilic phototrophic bacteria of the genus *Ectothiorhodospira*. *Eur. J. Biochem.*, 149: 135-139. DOI: 10.1111/j.1432-1033.1985.tb08903.x
- Garcia-Esteva, R., M. Argandona, M. Reina-Bueno, N. Capote and F. Iglesias-Guerra *et al.*, 2006. The ectD gene, which is involved in the synthesis of the compatible solute hydroxyectoine, is essential for thermoprotection of the halophilic bacterium *Chromohalobacter salexigens*. *J. Bacteriol.*, 188: 3774-3784. DOI: 10.1128/JB.00136-06
- Gilboa, H., M. Kogut, S. Chalamish, R. Regev and Y. Avi-Dor *et al.*, 1991. Use of ²³Na nuclear magnetic resonance spectroscopy to determine the true intracellular concentration of free sodium in a halophilic eubacterium. *J. Bacteriol.*, 173: 7021-7023. PMID: 1938904
- Goller, K., A. Ofer and E.A. Galinski, 1998. Construction and characterization of a NaCl-sensitive mutant of *Halomonas elongata* impaired in ectoine biosynthesis. *FEMS Microbiol. Lett.*, 161: 293-300. PMID: 9570121
- Graf, R., S. Anzali, J. Buenger, F. Pfluecker and H. Driller, 2008. The multifunctional role of ectoine as a natural cell protectant. *Clin. Dermatol.*, 26: 326-333. DOI: 10.1016/j.clindermatol.2008.01.002
- Grant, W.D., 2004. Life at low water activity. *Phil. Trans. R. Soc. Lond. B.*, 359: 1249-1267. DOI: 10.1098/rstb.2004.1502
- Hausinger, R.P., 2004. FeII/alpha-ketoglutarate-dependent hydroxylases and related enzymes. *Crit. Rev. Biochem. Mol. Biol.*, 39: 21-68. PMID: 15121720
- Hong, M., M. Fuangthong, J.D. Helmann and R.G. Brennan, 2005. Structure of an OhrR-ohrA operator complex reveals the DNA binding mechanism of the MarR family. *Mol. Cell*, 20: 131-141. DOI: 10.1016/j.molcel.2005.09.013
- Ikai, H. and S. Yamamoto, 1997. Identification and analysis of a gene encoding L-2,4-diaminobutyrate: 2-ketoglutarate 4-aminotransferase involved in 1,3-diaminopropane production pathway in *Acinetobacter baumannii*. *J. Bacteriol.*, 179: 5118-5125. PMID: 9260954

- Jebbar, M., R. Talibart, K. Gloux, T. Bernard and C. Blanco, 1992. Osmoprotection of *Escherichia coli* by ectoine: uptake and accumulation characteristics. J. Bacteriol., 174: 5027-5035. PMID: 1629159
- Kalyuzhnaya, M.G., V.N. Khmelenina, B.T. Eshinimaev, N.E. Suzina and D. Nikitin *et al.*, 2001. Taxonomic characterization of new alkaliphilic and alkalitolerant methanotrophs from soda lakes of the Southeastern Transbaikal region and description of *Methylomicrobium buryatense* sp. nov. Syst. Applied Microbiol., 24: 166-176. PMID: 11518319
- Kalyuzhnaya, M.G., V. Khmelenina, B. Eshinimaev, D. Sorokin and H. Fuse *et al.*, 2008. Classification of halo (alkali) philic and halo (alkali) tolerant methanotrophs provisionally assigned to the genera *Methylomicrobium* and *Methylobacter* and emended description of the genus *Methylomicrobium*. Int. J. Syst. Evol. Microbiol., 58: 591-596. DOI: 10.1099/ijms.0.65317-0
- Kempf, B. and E. Bremer, 1998. Uptake and synthesis of compatible solutes as microbial stress responses to high-osmolality environments. Arch. Microbiol., 170: 319-330. DOI: 10.1007/s002030050649
- Khmelenina, V.N., M.G. Kalyuzhnaya, V.G. Sakharovsky, N.E. Suzina and Y.A. Trotsenko *et al.*, 1999. Osmoadaptation in halophilic and alkaliphilic methanotrophs. Arch. Microbiol., 172: 321-329. DOI: 10.1007/s002030050786
- Kraegeloh, A. and H.J. Kunte, 2002. Novel insights into the role of potassium for osmoregulation in *Halomonas elongata*. Extremophiles, 6: 453-462. DOI: 10.1007/s00792-002-0277-4
- Kuhlmann, A.U. and E. Bremer, 2002. Osmotically regulated synthesis of the compatible solute ectoine in *Bacillus pasteurii* and related *Bacillus* spp. Applied Environ. Microbiol., 68: 772-783. DOI: 10.1128/AEM.68.2.772-783.2002.
- Kuhlmann, A.U., J. Bursy, S. Gimpel, T. Hoffmann and E. Bremer, 2008. Synthesis of the compatible solute ectoine in *Virgibacillus pantothenicus* is triggered by high salinity and low growth temperature. Applied Environ. Microbiol., 74: 4560-4563. DOI: 10.1128/AEM.00492-08
- Lo, C.C., A.C. Bonner, G. Xie, M.D. Souza and A.R. Jensen, 2009. Cohesion group approach for evolutionary analysis of aspartokinase, an enzyme that feeds a branched network of many biochemical pathways. Microbiol. Mol. Biol. Rev., 73: 594-651. DOI: 10.1128/MMBR.00024-09
- Louis, P. and E.A. Galinski, 1997. Characterization of genes for the biosynthesis of the compatible solute ectoine from *Marinococcus halophilus* and osmoregulated expression in *Escherichia coli*. Microbiology, 143: 1141-1149. DOI: 10.1099/00221287-143-4-1141
- Malin, G. and A. Lapidot, 1996. Induction of synthesis tetrahydropyrimidine derivatives in *Streptomyces* strain and their effect on *Escherichia coli* in response to osmotic and heat stress. J. Bacteriol., 178: 385-395. PMID: 8550457
- Marhuenda-Egea, F.C. and M.J. Bonete, 2002. Extreme halophilic enzymes in organic solvents. Curr. Opin. Biotechnol., 13: 385-389. DOI: 10.1016/S0958-1669(02)00338-5
- McDonough, M.A., K.L. Kavanagh, D. Butler, T. Searls and U. Oppermann 2005. Structure of human phytanoyl-CoA2-hydroxylase identifies molecular mechanisms of Refsum disease. J. Biol. Chem., 280: 41101-41110. DOI: 10.1074/jbc.M507528200
- Mevarech, M., F. Frolow and L.M. Gloss, 2000. Halophilic enzymes: Proteins with a grain of salt. Biophys. Chem., 86: 155-64. PMID: 11026680
- Murrell, J.C., I.R. McDonald and B. Gilbert, 2000. Regulation of expression of methane monooxygenases by copper ions. Trends Microbiol., 8: 221-225. DOI: 10.1074/jbc.M507528200
- Mustakhimov, I.I., O.N. Rozova, A.S. Reshetnikov, V.N. Khmelenina and J.C. Murrell *et al.*, 2008. Characterization of the recombinant diamino butyric acid acetyltransferase from *Methylophaga thalassica* and *Methylophaga alcalica*. FEMS Microbiol. Lett., 283: 91-96. DOI: 10.1111/j.1574-6968.2008.01156.x
- Mustakhimov, I.I., A.S. Reshetnikov, V.N. Khmelenina and Y.A. Trotsenko, 2009. EctR-a novel transcriptional regulator of ectoine biosynthesis genes in the haloalkaliphilic methylotrophic bacterium *Methylophaga alcalica*. Dokl. Biochem. Biophys., 429: 305-308. PMID: 20101826
- Mustakhimov, I.I., A.S. Reshetnikov, A.S. Glukhov, V.N. Khmelenina and M.G. Kalyuzhnaya *et al.*, 2010. Identification and characterization of EctR1, a new transcriptional regulator of the ectoine biosynthesis genes in the halotolerant methanotroph *Methylomicrobium alcaliphilum* 20Z. J. Bacteriol., 192: 410-417. DOI: 10.1128/JB.00553-09
- Nagata, S., K. Adachi, K. Shirai and H. Sano, 1995. ²³Na NMR spectroscopy of free Na⁺ in the halotolerant bacterium *Brevibacterium* sp. and *Escherichia coli*. Microbiology, 141: 729-736. DOI: 10.1099/13500872-141-3-729

- Ono, H., K. Sawada, N. Khunajakr, T. Toa and M. Yamamoto *et al.*, 1999. Characterization of biosynthetic enzymes for ectoine as a compatible solute in a moderately halophilic eubacterium, *Halomonas elongata*. J. Bacteriol., 181: 91-99. PMID 9864317
- Onraedt, A., C. de muynck, B. Walcarius, W. Soetaert and E. Vandamme, 2004. Ectoine accumulation in *Brevibacterium epidermis*. Biotechnol. Lett., 26: 1481-1485. DOI: 10.1023/B:BILE.0000044448.86907.e4
- Peters, R. and E.A. Galinski and H.G. Truper, 1990. The biosynthesis of ectoine. FEMS Microbiol. Lett., 71: 157-162. DOI: 10.1016/0378-1097(90)90049-V
- Prabhu, J., F. Schauwecker, N. Grammel, U. Keller and M. Bernhard, 2004. Functional expression of the ectoine hydroxylase gene (thpD) from *Streptomyces chrysomallus* in *Halomonas elongata*. Applied Environ. Microbiol., 70: 3130-3132. DOI: 10.1128/AEM.70.5.3130-3132.2004
- Rao, D.R., K. Hariharan and K.R. Vijayalakshmi, 1969. A study of the metabolism of L- α -diaminobutyric acid in a *Xanthomonas* species. J. Biochem., 114: 107-115. PMID: 4390206
- Reshetnikov, A.S., I.I. Mustakhimov, V.N. Khmelenina and Y.A. Trotsenko, 2005. Cloning, purification and characterization of the diaminobutyrate acetyltransferase from the halotolerant methanotroph *Methylomicrobium alcaliphilum* 20Z. Biochemistry, 70: 878-883. PMID: 16212543
- Reshetnikov, A.S., V.N. Khmelenina and Y.A. Trotsenko, 2006. Characterization of the ectoine biosynthesis genes of haloalkalotolerant obligate methanotroph "*Methylomicrobium alcaliphilum* 20Z". Arch. Microbiol., 184: 286-297. DOI: 10.1007/s00203-005-0042-z
- Reuter, K., M. Pittelkow, J. Bursy, A. Heine, T. Craan and E. Bremer, 2010. Synthesis of 5-hydroxyectoine from ectoine: Crystal structure of the non-heme Iron(II) and 2-oxoglutarate-dependent dioxygenase EctD. PLoS One, 5: e10647-e10647. DOI: 10.1371/journal.pone.0010647
- Roberts, M.F., 2004. Osmoadaptation and osmoregulation in archaea: Update. Front. Biosci., 9: 1999-2019. PMID: 15353266
- Roberts, M.F., 2005. Organic compatible solutes of halotolerant and halophilic microorganisms. Saline Syst., 1: 1-5. DOI: 10.1186/1746-1448-1-5
- Sauer, T. and E.A. Galinski, 1998. Bacterial milking a novel bioprocess for production of compatible solutes. Biotechnol. Bioeng., 57: 306-313. DOI: 10.1002/(SICI)1097-0290(19980205)57:3<306::AID-BIT7>3.0.CO;2-L
- Saum, S.H. and V. Muller, 2007. Salinity-dependent switching of osmolyte strategies in a moderately halophilic bacterium: glutamate induces proline biosynthesis in *Halobacillus halophilus*. J. Bacteriol., 189: 6968-6975. DOI: 10.1128/JB.00775-07
- Saum, S.H. and V. Muller, 2008a. Growth phase-dependent switch in osmolyte strategy in a moderate halophile: Ectoine is a minor osmolyte but major stationary phase solute in *Halobacillus halophilus*. Environ. Microbiol., 10: 716-726. DOI: 10.1111/j.1462-2920.2007.01494.x
- Saum, S.H. and V. Muller, 2008b. Regulation of osmoadaptation in the moderate halophile *Halobacillus halophilus*: chloride, glutamate and switching osmolyte strategies. Saline Syst., 4: 1-4. DOI: 10.1186/1746-1448-4-4
- Severin, J., A. Wohlfarth and E.A. Galinski, 1992. The predominant role of recently discovered tetrahydropyrimidines for the osmoadaptation of halophilic eubacteria. J. Gen. Microbiol., 138: 1629-1638. DOI: 10.1099/00221287-138-8-1629
- Toney, M.D., E. Hohenester, J.W. Keller and N. Jansonius, 1995. Structural and mechanistic analysis of two refined crystal structures of the pyridoxal phosphate-dependent enzyme dialkylglycine decarboxylase. J. Mol. Biol., 245: 151-179. DOI: 10.1006/jmbi.1994.0014
- Vargas, C., M. Argandona, M. Reina-Bueno, J. Rodriguez-Moya and C. Fernandez-Aunio *et al.*, 2008. Unraveling the adaptation responses to osmotic and temperature stress in *Chromohalobacter salexigens*, a bacterium with broad salinity tolerance. Saline Syst., 4: 1-14. DOI: 10.1186/1746-1448-4-14
- Ventosa, A. and J.J. Nieto, 1995. Biotechnological applications and potentialities of halophilic microorganisms. World J. Microbiol. Biotechnol., 11: 85-94. DOI: 10.1007/BF00339138
- Wilkinson, S.P. and A. Grove, 2006. Ligand-responsive transcriptional regulation by members of the MarR family of winged helix proteins. Curr. Issues Mol. Biol., 8: 51-62. PMID: 16450885
- Zhilina, T.N. and G.A. Zavarzin, 1990. Extremely halophilic, methylotrophic, anaerobic bacteria. FEMS Microbiol. Rev., 87: 315-322. DOI: 10.1111/j.1574-6968.1990.tb04930.x